

PERIOPERATIVE CARE MEASURES GROUP
 (For Patients Aged 18 and Older)
SuperBill Addendum

Patient Name: _____	Patient MRN: _____	Date of Service: _____	Physician: _____
------------------------	-----------------------	---------------------------	---------------------

CPT Code:

Measure Number and Title	Action Performed	Action Not Performed / Reason Documented	Action Not Performed / Reason Not Documented
20: Perioperative Care: Timing of Antibiotic Prophylaxis – Ordering Physician	<input type="checkbox"/> 4047F Documentation of order for prophylactic antibiotics to be given within one hour (if fluoroquinolone or vancomycin, two hours) prior to surgical incision (or start of procedure when no incision is required) <input type="checkbox"/> 4048F Documentation that prophylactic antibiotic was given within one hour (if fluoroquinolone or vancomycin, two hours) prior to surgical incision (or start of procedure when no incision is required)	<input type="checkbox"/> 4047F-1P Not given for medical reasons	<input type="checkbox"/> 4047F-8P Not given. No reason noted.
21: Perioperative Care: Selection of Prophylactic Antibiotic – First OR Second Generation Cephalosporin	<input type="checkbox"/> 4041F Documentation of order for cefazolin OR cefuroxime for antimicrobial prophylaxis	<input type="checkbox"/> 4041F-1P Not ordered for medical reasons	<input type="checkbox"/> 4041F-8P Not ordered. No reason noted.

<p>22: Perioperative Care: Discontinuation of Prophylactic Antibiotics (Non-Cardiac Procedures)</p>	<p><input type="checkbox"/> 4049F & 4046F Documentation that order was given to discontinue prophylactic antibiotics within 24 hours of surgical end time and documentation that prophylactic antibiotics were given within 4 hours prior to surgical incision or given intraoperatively</p>	<p><input type="checkbox"/> 4049F-1P & 4046F Documentation of medical reason(s) for not discontinuing prophylactic antibiotics within 24 hours of surgical end time and documentation that prophylactic antibiotics were given within 4 hours prior to surgical incision or given intraoperatively</p> <p><input type="checkbox"/> 4042F Documentation that prophylactic antibiotics were neither given within 4 hours prior to surgical incision nor given intraoperatively</p>	<p><input type="checkbox"/> 4049F-8P & 4046F Order was not given to discontinue prophylactic antibiotics within 24 hours of surgical end time, non-cardiac procedure, no reason noted and Documentation that prophylactic antibiotics were given within 4 hours prior to surgical incision or given intraoperatively</p>
<p>23: Perioperative Care: Venous Thromboembolism (VTE) Prophylaxis (When Indicated in ALL Patients)</p>	<p><input type="checkbox"/> 4044F Documentation that an order was given for venous thromboembolism (VTE) prophylaxis to be given within 24 hours prior to incision time or 24 hours after surgery end time.</p> <p>Use this option to note when VTE prophylaxis is ordered or VTE prophylaxis is given.</p>	<p><input type="checkbox"/> 4044F-1P VTE Prophylaxis not ordered for medical reasons</p>	<p><input type="checkbox"/> 4044F-8P VTE Prophylaxis not ordered. No reason noted.</p>